			CLASSIFIC	CATION GONFIDE	NETTAL /SECT	TRITTY THE	RMATTON		708
				ENTRAL INTELLI			OMMITON.		<u></u>
		x .	INF	ORMATIO	N REF	PORT			
· C	OUNTRY	•	USSR				DATE DISTR.	May	1953
S	UBJECT		Kyanite	e in the USSR	Ź		NO. OF PAGE	S l	50X1
	LACE CQUIRE	D					NO. OF ENCL	S.	50X1
	ATE CQUIRE	:D					SUPPLEMENT REPORT NO.	TO	
DA	TE OF	INFORMATION					*		50X1
	OF THE UNITE AND 794, OF LATION OF IT PROHIBITED B	O STATES, BITHIN THE M THE U.S. CODE, AS AMEN S CONTENTS TO OR RECE! Y LAW. THE REPROSUCT!	EANING OF TITLE 18, OED. ITS TRANSMIE FT BY AN UNAUTHOR! ON OF THIS FORM IS	SECTIONS 793 SSION ON REVE- IZED PERSON IS D PROHIBITED.	Т	HIS IS UNE	EVALUATED INF	ORMATI	NC
1.									
		Ac for so T							
		acondidation of a	Tr [atemena	from the mine	rals of th	ils group	isolation and - the polymon	Durc m	OUTLT
		considered s cations of t of them are very small a The deposits the form of They are cor	separately the Al203Si often four amounts, as of miners	the USSR kyani from the mine iO2 compounds and together, a s alumina high als of this gr lenses, strati metamorphic s	erals of the (andalusition) and all of refractor coup are of the (plant)	is group te, sillim them are ries (glin hydrothe	- the polymon anite and mul used, even th ozemnyye vyso rmal origin. enses and lay	rphic m lite). lough i bkoogne The o	All t be in upory). ccur in
2.		considered scations of to them are very small so the deposits	separately the Al203Si often four amounts, as of miners	from the mine iO2 compounds nd together, a salumina high als of this gr	erals of the (andalusition) and all of refractor coup are of the (plant)	is group te, sillim them are ries (glin hydrothe	- the polymon anite and mul used, even th ozemnyye vyso rmal origin. enses and lay	rphic m lite). lough i bkoogne The o	All t be in ipory). ccur in ilezhi) minantl
2.,		considered scations of them are very small as The deposits the form of They are cor Archaean). The Al ₂ O ₃ co within the scate 5/5/5/1	eparately the Al203Si often four mounts, as of miners pockets, i tained in ontent of the	from the mine iO2 compounds nd together, a salumina high als of this gr	rals of the (andalusity and all of refractor coup are of filed (place) schists and and other the coup are of the coup are of the coup are of the coup are of the coup and other the coup are of the coup are o	nis group te, sillim them are ries (glin f hydrothe stovyye) l d secondar minerals o nts are fo	- the polymoranite and mulused, even thousand origin. enses and lay quartzites f this group r an alumina int of as low	rphic milite). lite). lough i bkoogne The o rers (z (predo varies conten	All t be in ipory). ccur in alezhi) minantl 50X1 50X1 even t of at Al203 s 98%
2.		considered scations of to them are very small a The deposits the form of They are con Archaean). The Al ₂ O ₃ co within the sleast 54% in are used.	eparately the Al203Si often four mounts, as of minere pockets, trained in ontent of same depose these ore	from the mine iO2 compounds nd together, a s alumina high als of this gr lenses, strati metamorphic s the kyanites a it. Although es, in practic es the kyanite	rals of the (andalusity and all of a refractor roup are or iffied (plass schists and other requirements of content are content	nis group te, sillim them are ries (glin hydrothe stovyye) l i secondar minerals on ts are fo th a conte sometimes	- the polymorante and mulused, even those mal origin. enses and lay quartzites f this group r an aluminant of as low mounts to as	pnic m lite). lough i skoogne The o vers (z (predo varies conten as 45% much a	All t be in ipory). ccur in alezhi) minanti 50X1 50X1 even t of at Al203 s 98% 50X1 50X1
		considered scations of to five mare very small a the deposits the form of They are con Archaean). The Al ₂ O ₃ co within the sleast 54% in are used. I know of a northern spustation (station (station station station) is sovskoye and skiy Rayon is stations of the station stations station stations station stations stations station stations s	eparately the Al203Si often four mounts, as of minere pockets, intained in ontent of same deposit these ore in some ore deposits of the antsiya B. f deposits i Mikhaylo in Karel's	from the mine iO2 compounds and together, a s alumina high als of this gr lenses, strati metamorphic s the kyanites a it. Although es, in practic es the kyanite f kyanite in t Stanovoy Khre Never). This in Kochkarski vskoye deposit kaya ASSR _now	rals of the (andalusity and all of a refractor coup are or iffied (plans that and other requirement are content are content are the souther about the souther are that are the souther are that are the souther are the southe	nis group the, sillim them are ries (glin hydrothe stovyye) l isecondar minerals on the are fo the conte sometimes rn part of e), 350 ki o-called on n Chelyabi posits on	- the polymora anite and mulused, even those origin. enses and lay y quartzites of this group ran aluminant of as low mounts to as 'Yakutskaya'. Inskaya Oblas the Kem Rive:	pnic m lite). lite). lough i koogne The o rers (z (predo varies conten as 45% much a ASSR, i n Bel's del's tt - the r in Sh	All t be in lipory). ccur in alezhi) minantl; 50X1 even t of at Al203 s 98% 50X1 50X1 n the hoy Nev . I Bori- uyeret-
		considered scations of to fine are very small a The deposits the form of They are con Archaean). The Al ₂ O ₃ co within the sleast 54% in are used. I know of a northern spi station (staken are consequence of the consequence of the consequence of the controllers are consequence of the consequenc	eparately the Al203Si often four mounts, as of minere pockets, intained in ontent of same deposit these ore in some ore deposits of the antsiya B. f deposits i Mikhaylo in Karel's	from the mine iO2 compounds not together, a salumina high als of this grienses, strati metamorphic stratically and the kyanites a it. Although es, in practices the kyanite f kyanite in together in Kochkarski vskoye deposit kaya ASSR /nowee.	rals of the (andalusity and all of a refractor coup are or iffied (plans that and other requirement are content are content are the souther about the souther are that are the souther are that are the souther are the southe	nis group the, sillim them are ries (glin hydrothe stovyye) l isecondar minerals on the are fo the conte sometimes rn part of e), 350 ki o-called on n Chelyabi posits on	- the polymora anite and mulused, even those origin. enses and lay y quartzites of this group ran aluminant of as low mounts to as 'Yakutskaya'. Inskaya Oblas the Kem Rive:	pnic m lite). lite). lough i koogne The o rers (z (predo varies conten as 45% much a ASSR, i n Bel's del's tt - the r in Sh	All t be in lipory). ccur in alezhi) minanti 50X1 even t of at Al203 s 98% 50X1 50X1 the hoy Nev . Bori- uyeret-
		considered scations of to five mare very small a the deposits the form of They are con Archaean). The Al ₂ O ₃ co within the sleast 54% in are used. I know of a northern spustation (station (station station station) is sovskoye and skiy Rayon is stations of the station stations station stations station stations stations station stations s	eparately the Al203Si often four mounts, as of minere pockets, intained in ontent of same deposit these ore in some ore deposits of the antsiya B. f deposits i Mikhaylo in Karel's	from the mine iO2 compounds not together, a salumina high als of this grienses, strati metamorphic stratically and the kyanites a it. Although es, in practices the kyanite f kyanite in together in Kochkarski vskoye deposit kaya ASSR /nowee.	rals of ti (andalusi) (andalusi) (and all of a refractor (oup are or lified (place (place) (pl	nis group the, sillim them are ries (glin hydrothe stovyye) l isecondar minerals on the are fo the conte sometimes rn part of e), 350 ki o-called on n Chelyabi posits on	- the polymora anite and mulused, even those origin. enses and lay y quartzites of this group ran aluminant of as low mounts to as 'Yakutskaya'. Inskaya Oblas the Kem Rive:	pnic m lite). lite). lough i koogne The o rers (z (predo varies conten as 45% much a ASSR, i n Bel's del's tt - the r in Sh	All t be in lipory). ccur in alezhi) minantl; 50X1 50X1 even t of at Al203 s 98% 50X1 50X1 n the hoy Nev Bori- uyeret- some
		considered scations of to five mare very small a the deposits the form of They are con Archaean). The Al ₂ O ₃ co within the sleast 54% in are used. I know of a northern spustation (station (station station station) is sovskoye and skiy Rayon is stations of the station stations station stations station stations stations station stations s	eparately the Al203Si often four mounts, as of minere pockets, intained in ontent of same deposit these ore in some ore deposits of the antsiya B. f deposits i Mikhaylo in Karel's	from the mine iO2 compounds not together, a salumina high als of this grienses, strati metamorphic stratically and the kyanites a it. Although es, in practices the kyanite f kyanite in together in Kochkarski vskoye deposit kaya ASSR /nowee.	rals of ti (andalusi) (andalusi) (and all of a refractor (oup are or lified (place (place) (pl	nis group the, sillim them are ries (glin hydrothe stovyye) l isecondar minerals on the are fo the conte sometimes rn part of e), 350 ki o-called on n Chelyabi posits on	- the polymora anite and mulused, even those origin. enses and lay y quartzites of this group ran aluminant of as low mounts to as 'Yakutskaya'. Inskaya Oblas the Kem Rive:	pnic m lite). lite). lough i koogne The o rers (z (predo varies conten as 45% much a ASSR, i n Bel's del's tt - the r in Sh	All t be in lipory). ccur in alezhi) minant! 50X1 50X1 even t of at Al203 s 98% 50X1 50X1 n the hoy Nev Bori- uyeret- some
		considered scations of to five mare very small a the deposits the form of They are con Archaean). The Al ₂ O ₃ co within the sleast 54% in are used. I know of a northern spustation (station (station station station) is sovskoye and skiy Rayon is stations of the station stations station stations station stations stations station stations s	eparately the Al203Si often four mounts, as of minere pockets, intained in ontent of same deposit these ore in some ore deposits of the antsiya B. f deposits i Mikhaylo in Karel's	from the mine iO2 compounds not together, a salumina high als of this grienses, strati metamorphic stratically and the kyanites a it. Although es, in practices the kyanite f kyanite in together in Kochkarski vskoye deposit kaya ASSR /nowee.	rals of ti (andalusi) (andalusi) (and all of a refractor (oup are or lified (place (place) (pl	nis group the, sillim them are ries (glin hydrothe stovyye) l isecondar minerals on the are fo the conte sometimes rn part of e), 350 ki o-called on n Chelyabi posits on	- the polymora anite and mulused, even those origin. enses and lay y quartzites of this group ran aluminant of as low mounts to as 'Yakutskaya'. Inskaya Oblas the Kem Rive:	pnic m lite). lite). lough i koogne The o rers (z (predo varies conten as 45% much a ASSR, i n Bel's del's tt - the r in Sh	All t be in lipory). ccur in alezhi) minant! 50X1 50X1 even t of at Al203 s 98% 50X1 50X1 n the hoy Nev Bori- uyeret- some
		considered scations of to five mare very small a the deposits the form of They are con Archaean). The Al ₂ O ₃ co within the sleast 54% in are used. I know of a northern spustation (station (station station station) is sovskoye and skiy Rayon is stations of the station stations station stations station stations stations station stations s	eparately the Al203Si often four mounts, as of minere pockets, intained in ontent of same deposit these ore in some ore deposits of the antsiya B. f deposits i Mikhaylo in Karel's	from the mine iO2 compounds not together, a salumina high als of this grienses, strati metamorphic stratically and the kyanites a it. Although es, in practices the kyanite f kyanite in together in Kochkarski vskoye deposit kaya ASSR /nowee.	rals of ti (andalusi) (andalusi) (and all of a refractor (oup are or lified (place (place) (pl	nis group the, sillim them are ries (glin hydrothe stovyye) l isecondar minerals on the are fo the conte sometimes rn part of e), 350 ki o-called on n Chelyabi posits on	- the polymora anite and mulused, even those origin. enses and lay y quartzites of this group ran aluminant of as low mounts to as 'Yakutskaya'. Inskaya Oblas the Kem Rive:	pnic m lite). lite). lough i koogne The o rers (z (predo varies conten as 45% much a ASSR, i n Bel's del's tt - the r in Sh	All t be in lipory). ccur in alezhi) minantl; 50X1 50X1 even t of at Al203 s 98% 50X1 50X1 n the hoy Nev Bori- uyeret- some
		considered scations of to them are very small a The deposits the form of They are con Archaean). The Al ₂ O ₃ co within the sleast 54% in are used. I know of a northern spi station (steknow also or sovskoye and skiy Rayon deposits of	eparately the Al203Si often four mounts, as of minere pockets, intained in ontent of same deposit these ore in some ore deposits of the antsiya B. f deposits i Mikhaylo in Karel's	from the mine iO2 compounds nod together, a salumina high als of this grienses, strati metamorphic stranding the kyanites a it. Although es, in practices the kyanite in Kochkarski vskoye deposit kaya ASSR /now e.	and other and ot	nis group the, sillim them are ries (glin f hydrothe stovyye) l d secondar minerals o nts are fo th a conte sometimes rn part of e), 350 ki o called o n Chelyabi posits on innish SSF	- the polymoranite and mulused, even thoused, even thousens and lay conservations of this group ran alumina and of as low mounts to as a lowesters from the constant of the co	pnic m lite). lite). lough i koogne The o rers (z (predo varies conten as 45% much a ASSR, i n Bel's del's tt - the r in Sh	All t be in lipory). ccur in alezhi) minant! 50X1 50X1 even t of at Al203 s 98% 50X1 50X1 n the hoy Nev Bori- uyeret- some